After Your Surgery

Closed Reduction of Nasal Fracture

CAPITAL ENT & SINUS CENTER

What to Expect

- Mild bleeding from the nose for the first few days.
- You may have soft plastic splints inside your nose held in place by a stitch. These are removed at your first post-op visit.
- Nasal pain and congestion are normal. You may feel more congested than you did before surgery because of swelling and the splints (if in place).
- Your upper front teeth may feel numb. This almost always resolves within a few weeks.
- Your nose is fragile for at least 6 weeks after surgery. Avoid any injury to the nose during this period.
- You may have a splint on the outside of your nose, held in place with tape and surgical glue. It is removed at your first post-op visit.

What to Do

- Use nasal saline spray, 2 sprays in each nostril every 2 to 3 hours while awake (5 or 6 times a day), until your follow-up visit.
- You may use Afrin nasal spray as needed for bleeding and nasal congestion for 3 to 5 days after surgery. Stop Afrin after 3 to 5 days.
- For the first 3 nights, sleep with your head and shoulders raised 30 to 45 degrees (wedge, pillows or recliner).
- Cold pack in a thin towel: 15 to 20 minutes on, 20 off, while awake for the first 48 hours; do not press on the nose.
- Tape a piece of gauze under your nose if there is mild bleeding. Change the gauze as needed.
- Tylenol 1,000 mg every 6 hours (max 4,000 mg a day; 3,000 mg if the package says so or with liver disease) and ibuprofen 400 mg every 8 hours with food, unless your surgeon said not to or you have an ulcer or kidney disease.
- If the external splint falls off early, gently place it back on the nose and secure it with medical tape from a pharmacy.

What Not to Do

- After sedation or general anesthesia: for 24 hours, no driving, alcohol or important decisions; after general anesthesia, we recommend an adult stay with you the first night.
- No heavy lifting (over about 10 lb) or strenuous exercise (anything that makes you breathe hard, strain or sweat) for 2 weeks after surgery. Light jogging is fine.
- Do not blow your nose for the first week. Do not sneeze with your mouth closed.
- Contact sports: usually no sooner than 3 weeks, with a protective face mask until at least 6 weeks. Combat sports such as boxing: usually about 3 months. Check with us before returning.

When to Call or See the Doctor

- Temperature of 101° F+ for over 24 hours or starting 3+ days after the procedure, or 102° F+ at any time (check before the next Tylenol or ibuprofen dose).
- Nasal blockage, pressure or pain inside the nose that gets worse in the first few days instead of better: call us promptly.
- Worsening, severe nasal pain.
- Bleeding from the nose that does not stop, or any bleeding that worries you: call us for guidance. Severe bleeding you cannot control, or bleeding with lightheadedness or fainting: call 911 or go to the nearest emergency room.

Questions or concerns? Call the office that scheduled your surgery. After hours, call our main number, 512-339-4040, to reach the on-call physician. For a life-threatening emergency, call 911.

Austin 512-339-4040 **Lakeway** 512-682-4798 **Marble Falls** 830-992-2830

Dripping Springs 512-339-4040

Scan for the full guide